



Dairy Stockmanship – Empowering People and Cattle Humane Livestock Handling: Practical Applications of Animal Behavior and Welfare Science

North Carolina State University - College of Veterinary Medicine Outreach

Don Höglund MS, DVM

djhoglund@ncsu.edu

Introduction

The veterinary profession and dairy producers have taken a strong interest in dairy stockmanship – the interactions that occur between people and cows. The daily management of a dairy operation requires personnel to interact with the cattle many times a day. Dairy stockmanship is the implementation of low-energy cow handling techniques to improve the outcomes for both the people and the cattle.

Dairy Stockmanship

In stockmanship terms stimulation of any kind on livestock is often referred to as a form of pressure. Behaviorists prefer to use the word stimulus instead of mental pressure. Stimulus can be quantified whereas assessments of mental pressure are hard to measure. In discussions on stockmanship or cattle handling people frequently refer to flight or escape zones of an animal as an arbitrary measure of how much stimulus or encroachment an animal will endure prior to fleeing or fighting. Conceptually, the flight or escape zone can be thought of as being an animal's personal space and when that space is violated the animal may determine that it is no longer safe and react. In practical terms, as a human approaches livestock the animal begins to feel stimulus from human encroachment. The exact flight distance and the extent of the response to human presence may vary animal to animal or within the same animal depending on the various factors influencing the animal, such as, prior animal experiences, previous human interaction, the distance between the human and animal, natural or artificial boundaries, husbandry practices, age of the animal, other competing environment stimuli, health and well-being of the animal(s) in general, and even the time of day. By closely observing the response of the animal approached, livestock handlers can be able to observe and learn from the effects of the stimulus on animal behavior. Alternatively, by moving away from animals some or all of the stimulus may be relieved and understanding this animal behavior will also be useful to livestock handlers. The key point is that handlers can induce animals to move, turn, or stop by exerting and manipulating stimulus.

Every interaction between people and cows shapes the future behavior of both. These interactions can be positive or negative but are very rarely neutral. The concept of stockmanship, or low-energy handling of livestock helps people become aware of human behavior and the impact it has on livestock. With proper handling cattle are easier to work and move and that creates a desirable environment for both cattle and people. In the dairy industry people interact with cows several times a day and these situations present opportunities to create positive, human and animal interactions.

General Stockmanship Concepts

There are a few general concepts about cow behavior stock handlers should keep in mind. Livestock derive information from the environment through their five senses: sight, hearing, smell, touch and taste. Cows do not use language to communicate with people so stock handlers must communicate with cows by stimulating the senses of the animal. The two most important senses a cow uses to understand what is going on in her environment are sight and hearing.

Cows consistently look at what is stimulating them. Because the eyes and ears of the cow are positioned on the side of the skull, cows have excellent peripheral vision and hearing. There is a narrow blind spot directly behind her rump. A good general rule is that if the handler can see the cow's eyeball she can probably see the handler. Therefore, the human should approach the animal from a position where her eyeball can be seen, in this manner she can probably see and hear the handler. Surprising livestock is never a good idea, so let them see the handler if possible and if not, let them gently hear who approaches them.

Cows tend to move in an arc around whatever they perceive as stimulus. This allows them to keep an eye on what is stimulating them as they move around or away from it. Cows tend to follow other cows. These two concepts are invaluable when emptying a cattle pen or loading a transport with cattle. If the handler can create positive motion at the front of the herd and then avoid doing anything to slow or stop the flow, cows will tend to move in the direction they are facing while following the cow in front of them. If, for example, handlers are moving animals into the parlor, the task will be accomplished more efficiently if the handler induces the animals to face the opening into the parlor. If the handler causes the animals to turn back toward the crowd gate, flow stops and the cattle tend to bunch. Handlers need to pay close attention to their position in relationship to the direction of cow movement. It is most important not to over-stimulate or to apply stimulus in an unpredictable manner to the animal. Extreme examples of over-stimulation are shouting, arm waving, and hitting animals or using electric prods to get them to move. Cows do not respond positively when over-stimulated, they exhibit agitation and may run potentially leading to harm. These examples of too much stimulus can be called high-energy cow handling techniques.

Herding a cow properly involves the right approach angle, speed, and timing. There is no complicated or magic formula. The cow's behavior will inform the handler if the angle, speed,

and timing were correct. If she didn't respond as the handler intended, then the handler should back along the same line as the approach, change the angle and the speed of approach. However, one concept has universal importance in moving cattle and it is that driving cattle from directly behind them, in their blind spot, causes the animal to turn and face the handler in order to get at least one eye on the stimulus. That handling mistake stops the forward motion of cattle because a cow tends not to walk far with her head turned. Cows seem to follow their eyes.

Cows walk at about two-miles per hour (mph) while people tend to walk about three to four mph. Handlers walking at their normal pace and parallel with cows will eventually overtake the cow, first slowing them and then stopping forward motion altogether. Handlers need to recognize this and slow their walking speed in order to move at the same pace as calm cows. Since it generally takes more stimulus to start a cow moving than it does to keep her moving, once cow motion begins the handler should slow or pause momentarily in order to create some distance between themselves and the moving cow. The handler then continues to apply only the stimulus needed to keep the cow moving calmly. Over-stimulating in order to start motion or during movement frequently causes cows to over-react and run. This is often seen when moving heifers.

Walking parallel against the flow of cows tends to speed them. This works because cows want to go the direction they are facing and they want to get away from the human stimulus; especially the human face and eyes. Walking parallel against the flow of cattle can help load or unload a chute, transport, or parlor, and is valuable when encouraging cows to exit the return alley. If more than one person is in the vicinity of the same animal or group of animals, it is best that one person stimulate at a time. With two or more handlers, it is very easy to apply conflicting stimulus to the cattle. Understandably, this would result in conflicting stimuli to the cows and results in poor communication to the animal. Consistent handling methods allow the cows to know what will happen next and that seems to have a calming effect on herd animals.

A good time to work animals is when they first arrive to a new pen or facility. Examples of this are during weaning of heifers from hutches into group pens or immediately after springing heifers arrive at a facility new to them. Spending 10-20 minutes allows handlers to develop a calm relationship with the new cattle while introducing the animals to the new environment. This also creates a great opportunity to examine those animals for any health problems.

As people learn to apply stockmanship skills on cattle operations a frequent question arises about what to do with new cattle? As simple as it may sound, the answer is that the behavior of the animals will tell handlers what should done with them. For example, if cattle run, back and forth or circle non-stop, the handlers need to slow that motion. If the cows bunch in a corner and have no movement, a handler or at most a few handlers should create slow movement that involves teaching new cattle to accept human stimulus. This also helps animals learn the boundaries of their new confinement while teaching them where food and water exists. Each

time cattle are worked properly they learn and become easier to work the next time. That is to say that animals learn calm handling if handled calmly. When livestock operations only consider working cattle if specific tasks are to be accomplished (such as vaccinating) a negative impression of handling can be imprinted in the cows' memory. Naturally, negative interactions can make cows become harder to handle over time. Frequently, we find that the older cows in a herd can be difficult to move. We must understand that their current behavior is the sum total of the interactions with humans over her lifetime; positive and negative. Dairy stockmanship is about reconnecting people with dairy cows for positive outcomes and it is fundamentally about learning how cows respond to the behaviors of people in a dynamic environment.

The Concept of Stress in Dairy Cattle

The general concept of low-stress handling is being widely discussed in the dairy industry today. The National Dairy FARM Program: Farmers Assuring Responsible ManagementSM, created by the National Milk Producers Federation (NMPF) with support from Dairy Management Inc. (DMI), specifically states as a best management practice "Employees should be properly trained to handle animals with a minimum of stress to the animal, and the consequences of inhumane handling should be known and enforced." The National Dairy FARM Program is designed to demonstrate that U.S. milk producers are committed to providing the highest standards of animal care and quality assurance. This voluntary program, available to all producers, provides a consistent on-farm animal well-being program that includes education, on-farm evaluations and third-party verification. Whether it be dairy stockmanship training or a program like FARM, the increased usage of the term "low-stress cattle handling techniques" has raised the questions of what exactly is stress, and how do we determine if it is "low" or "high"?

If you ask twelve people to define "stress" you would likely get 12 different answers. This creates an interesting challenge for us if we are going to attempt to determine the level of animal stress on a particular farm and whether the stress level is "low" or "high". If we struggle to define stress, how can we measure it? One of the goals of this paper is to introduce the reader to the scientific study of stress biology and to suggest that farm managers and advisors can utilize this understanding to assist in the evaluation whether cow handling stress is "low" or "high" on a dairy operation. A lengthy list of references is provided at the end of the paper for those interested in researching further into the concepts of animal stress biology.

A brief history of stress research pioneers will be helpful to understand how the term came into such widespread use. Hans Seyle (1907-1982) is generally recognized for being the first researcher to demonstrate the existence of biological stress. In 1936 Seyle defined stress as "the non-specific response of the body to any demand for change." Seyle demonstrated in his research that a wide variety of noxious stimuli caused a very consistent set of pathologic

changes in laboratory rats. Seyle's work created much interest and discussion in the scientific community.

The work of Robert Sapolsky is also useful in understanding the concept of biological stress. Sapolsky suggests a very useful approach by differentiating a "stressor" from the body's "stress response". Sapolsky defined a stressor as anything that disrupts physiological balance. A stress response is defined as the body's adaptations designed to re-establish the balance. Discussions at the 2011 Trends in Stress Biology course taught at Aarhus University suggested some slight refinements to the definitions.

- Stressor = event threatening or potentially threatening the homeostatic balance
- Stress Response = the body's attempt to re-establish the homeostasis after encountering a stressor.

Stressors can be described by their characteristics such as: duration, frequency, intensity, predictability, and ability to be controlled. It is important to note that while stressors can be physical things (heat, cold, starvation, etc.) psychological factors can also trigger the stress response in an animal in the absence of anything physically threatening to an animal.

Sapolsky in his writings proposes that the stress response evolved as adaptive survival mechanism for animals. It is now increasingly recognized that the consequences of the stress response can be maladaptive and that there is a "biological cost" to the animal for mounting a stress response. It is actually incorrect to state that stress makes an animal sick. To be correct, one should state that the stress response makes you more likely to get diseases that make you sick.

There is no single litmus test for stress because of the multiple ways the body responds to stressors. Since stressors will result in both behavioral responses and physiological responses on the part of the animal proper assessment of an animal's stress response requires one look both. One cannot interpret physiological test results without knowing the behavior.

An understanding of stockmanship principles will help one to be aware of behavior responses in animals. The physiological components of the stress response are significantly influenced by the endocrine system. Broadly speaking, all stressors provoke some degree of cortisol secretion as well as a multitude of other physiologic responses. The exact orchestration of the many hormones involved will vary depending on the stressor. In this way, different stressors have a different "stress signature" that describes the overall stress response. Work in this area is very interesting and in the future will most certainly allow us to improve and refine our evaluation of the physiological response to stress.

It is still our present understanding that glucocorticoids (cortisol) and catecholamine's (adrenalin) together mediate most of the changes that form the stress response. Today,

measuring cortisol remains the gold standard to evaluate the physiologic response to stressors. Researchers are actively engaged in searching for additional physiologic measures, but it is clear that cortisol does play an important role. Understanding the Hypothalamic-Pituitary-Adrenal axis (HPA axis) is critical to understanding the physiology of the stress response.

Blood sampling has been the traditional measure used to evaluate the cortisol level in an animal. However, plasma cortisol evaluation is not without issues. For example, obtaining a blood sample in itself can be stressful, especially in wildlife or zoo animals. Dr. Rupert Palme (Dept. Biomed. Sciences/Biochemistry, University of Veterinary Medicine, Vienna) and other researchers have been actively looking into alternatives to blood sampling. Cortisol is metabolized in the liver and cortisol metabolites are excreted in the urine and feces. Measuring cortisol metabolites in the feces (FCM's) has received a significant amount of attention. Since 1997, over 130 publications have used the measurement of FCM's on a wide variety of animal species, including dairy cattle.

The positive impact of better cow handling has been clearly demonstrated by Australia's Animal Welfare Science Centre, a joint organization with Australia's University of Melbourne, Monash University and the Victorian State Department of Primary Industries. The Centre is internationally recognized as a leading research and educational facility of animal welfare topics. Interested readers are directed to the Hemsworth references included in this paper for more detailed information.

Dairy veterinarians are frequently involved in on-farm training programs for dairy owners and their employees. Delivering effective training programs for dairy workers is a very valuable production medicine service to offer to dairy clients.

Our long-term goal is to develop useful training resources that dairy veterinarians will be able to utilize to improve their own stockmanship skills as well as use to facilitate on-farm training with dairy clients. In addition, we are actively developing a research program to study the cow behavior responses to stockmanship techniques.

References

Brown RE. 1994. An introduction to neuroendocrinology, chapters 1-4, Cambridge University Press, p. 1-55.

Cohen S, Kessler R, Gordon LU. *Measuring Stress: A Guide for Health and Social Scientists.* (1997, Oxford University Press) ISBN 978-0-19-512120-9

- Hansen SW, Malmkvist J, Palme R, Damgaard BM. 2007. Do double cages and access to occupational material improve the welfare of farmed mink? *Anim. Welf.* 16, 63-76
- Hayward LS, Wingfield (2004). Maternal corticosterone is transferred to avian yolk and may alter offspring growth and adult phenotype. *Gen. Comp. Endocrinol.* 135, 365-371.
- Hemsworth, P.H., Barnett, J.L., Tilbrook, A.J., Hansen, C., 1989. The Effects of Handling by Humans at Calving and During Milking on the Behaviour and Milk Cortisol Concentrations of Primiparous Dairy Cows. *Appl. Anim. Behav. Sci.* 22, 313-326.
- Hemsworth, P.H., Coleman, G.J., Barnett, J.L., Borg, S., 2000. Relationships between human-animal interactions and productivity of commercial dairy cows. *J. Anim. Sci.* 78, 2821-2831.
- Hemsworth, P.H., 2007. Ethical stockmanship. *Aust. Vet. J.* 85, 194-200.
- Herskin MS, Jensen KH. 2002. Effects of open field testing and associated handling v. handling alone on the adrenocortical reactivity of piglets around weaning. *J. Anim. Sci.* 74, 485-491.
- Jarvis S, Moinard C, Robson SK, Baxter E, Ormandy E, Douglas AJ, Seckl JR, Russell JA, Lawrence AB. 2006. Programming the offspring of the pig by prenatal social stress: Neuroendocrine activity and behaviour. *Horm. Behav.* 49: 68-80.
- Jensen P, Toates FM. 1997. Stress as a state of motivational systems. *Appl. Anim. Behav. Sci.* 53, 145-156.
- Jensen P, Andersson L. 2005. Genomics meets ethology: A new route to understanding domestication, behavior, and sustainability in animal breeding. *Ambio* 34, 320-324.
- Joëls M, Karst H, DeRijk R, de Kloet ER. 2007. The coming out of the brain mineralocorticoid receptor. *Trends Neurosci.* 31, 1-7.
- Laviola G, Hannan AJ, Macrì S, Solinas M, Jaber M. 2008. Effects of enriched environment on animal models of neurodegenerative diseases and psychiatric disorders. *Neurobiol. Disease* 31, 159-168.
- Macrì S, Mason GJ, Würbel H. 2004. Dissociation in the effects of neonatal maternal separations on maternal care and the offspring's HPA and fear responses in rats. *Eur. J. Neurosci.* 20, 1017-1024.
- Macrì S, Pasquali P, Bonsignore LC, Pieretti S, Cirulli F, Chiarotti F, Laviola G. 2007. Moderate neonatal stress decreases within-group variation in behavioural, immune and HPA responses in adult mice. *PLoS ONE* 2(10): e1015. doi:10.1371/journal.pone.0001015., 1-9
- Macrì S, Würbel H. 2006. Developmental plasticity of HPA and fear responses in rats: A critical review of the maternal mediation hypothesis. *Horm. Behav.* 50, 667-680.

- Malmkvist J, Damgaard BM, Pedersen LJ, Jørgensen E, Thodberg K, Chaloupkova H, Bruckmaier, RM. 2009. Effects of thermal environment on HPA-axis hormones, oxytocin and behavioral activity in periparturient sows. *J. Anim. Sci.* 87, 2796-2805.
- Malmkvist J, Palme R. 2008. Periparturient nest building: implications for parturition, kit survival, maternal stress and behaviour in farmed mink (*Mustela vison*). *Appl. Anim. Behav. Sci.* 114, 270-283.
- Mateo JM, Cavigelli SA. 2005. A validation for extraction methods for noninvasive sampling of glucocorticoids in free-living ground squirrels. *Physiol. Biochem. Zool.* 78, 1069-1084.
- Mellor DJ, Cook CJ, Stafford KJ. 2000. Quantifying some responses to pain as a stressor. Chap. 9 in GP Moberg and JA Mench (eds.), *The biology of animal stress*, CAB international, 171-198.
- Herskin MS, Munksgaard L, Ladewig J. 2004. Effects of acute stressors on nociception, adrenocortical responses and behaviour of dairy cows. *Physiol. Behav.* 83, 411-420.
- Heistermann M, Palme R, Ganswindt A. *Comparison of Different Enzymeimmunoassays for Assessment of Adrenocortical Activity in Primates Based on Fecal Analysis*. American Journal of Primatology 68:257–273 (2006)
- Moberg GP. 2000. Biological response to stress: Implications for animal welfare. Chapter 1 in GP Moberg and JA Mench (eds), *The biology of Animal Stress*, CAB international, p. 1-21.
- Mormede P, Andanson S, Auperin B, Beerda B, Guemene D, Malmkvist J, Manteca X, Manteuffel G, Prunet P, van Reenen CG, Richard S, Veissier I. 2007. Exploration of the hypothalamic-pituitaryadrenal function as a tool to evaluate animal welfare. *Physiol. Beh.* 92, 317-339.
- Morrow CJ, Kolver ES, Verkerk GA, Matthews LR. *Fecal Glucocorticoid Metabolites as a Measure of Adrenal Activity in Dairy Cattle*. General and Comparative Endocrinology 126, 229-241 (2002)
- Möstl E, Rettenbacher S, Palme R. 2005. Measurement of corticosterone metabolites in birds' droppings: An analytical approach. *Annals New York Acad. Sci.* 1046, 17-34.
- Palme R. 2005. Measuring fecal steroids: Guidelines for practical application. *Annals New York Acad. Sci.* 1046, 75-80.
- Palme R, Rettenbacher S, Touma C, El-Bahr SM, Möstl E. 2005. Stress hormones in mammals and birds: Comparative aspects regarding metabolism, excretion and noninvasive measurement in fecal samples. *Trends in Comparative Endocrinology and Neurobiology*. *Annals New York Acad. Sci.* 1040, 162-171.
- Palme R, Robia C, Baumgartner W, Mostl E. *Transport stress in cattle as reflected by an increase in faecal cortisol metabolite concentrations*. *Veterinary Record* (2000) 146, 108-109

Parker KJ, Maestripieri D. 2010. Identifying key features of early stressful experiences that produce stress vulnerability and resilience in primates. *Neurosci. Biobehav. Rev.* In press, doi: 10.1016/j.neubiorev.2010.09.003. 18pp

Sapolsky RM. 2002. Endocrinology of the Stress-Response. Chapter 11 in J. B. Becker, M. Breedlove, D. Crews, M. M. McCarty, *Behavioral Endocrinology*. 2nd edition. MIT Press, London, UK.

Sapolsky, R. *Why Zebras Don't Get Ulcers* (1994, Holt/Owl 3rd Rep. Ed. 2004) ISBN 978-0-8050-7369-0

Touma C, Palme R. 2005. Measuring fecal glucocorticoid metabolites in mammals and birds: The importance of validation. *Annals New York Acad. Sci.* 1046, 54-74.

Selye H. 1936. A syndrome produced by diverse nocuous agents. *Nature* 138, 32.

Weis JM. 1968. Effects of coping responses on stress. *J. Comp. Physiol. Psychol.* 65, 251-260.

Weis JM. 1971. Effect of coping behaviour in different warning signal conditions on stress pathology in rats. *J. Comp. Physiol. Psychol.* 77, 1-13.

Wolfer DP, Litvin O, Morf S, Nitsch RM, Lipp H-P, Würbel H. 2004. Cage enrichment and mouse behaviour. *Nature* 432, 821-822

NOTES